A Survey From Your Healthcare Provider — PHQ-9 Modified for Teens

Name		Clinician			
Medical Record or ID Number	Date				
Instructions: How often have you been bothered by eacl For each symptom put an "X" in the box beneath the ans					
	(O) Not At All	(1) Several Days	(2) More Than Half the Days	(3) Nearly Every Day	
Feeling down, depressed, irritable, or hopeless?					
2. Little interest or pleasure in doing things?					
3. Trouble falling asleep, staying asleep, or sleeping too much?					
4. Poor appetite, weight loss, or overeating?					
5. Feeling tired, or having little energy?					
6. Feeling bad about yourself — or feeling that you are a failure, or that you have let yourself or your family down?					
7. Trouble concentrating on things like school work, reading, or watching TV?					
8. Moving or speaking so slowly that other people could have noticed? Or the opposite — being so fidgety or restless that you were moving around a lot more than usual?					
Thoughts that you would be better off dead, or of hurting yourself in some way?					
10. In the <i>past year</i> have you felt depressed or sad most days, even it	f you felt okay so	metimes?	Yes N	o	
11. If you are experiencing any of the problems on this form, how difficult at all Somewhat difficult Very		oroblems made it for Extremely difficult	you to do your work,		
12. Has there been a time in the past month when you have had serio	us thoughts abou	it ending your life?	Yes N	o	
13. Have you ever, in your whole life, tried to kill yourself or made a si	uicide attempt?		Yes N	0	
		FOR OFFICE USE	ONLY Score		

Q. 12 and Q. 13 = Y or TS = ≥11

Administering, Scoring and Interpreting the PHQ-9 Screening Questionnaire

Administering

- The Patient Health Questionnaire Modified for Teens (PHQ-Modified) can be used with patients between the ages of 12 and 18 and takes less than five minutes to complete and score.
- The PHQ-9 Modified can be administered and scored by a nurse, medical technician, physical assistant, physician or other office staff.
- Patients should be left alone to complete the PHQ-9 Modified in a private area, such as an exam room or private area of the waiting room.
- Patients should be informed of their confidentiality rights before the PHQ-9 Modified is administered.
- The American Academy of Pediatrics and U.S. Preventive Service Task Force recommend that depression screening be conducted annually.

Scoring

For every X:

Not all =0 Several days=1

More than half the days=2

Nearly every day=3

Add up all "X" ed boxes on the screen

Defining a Positive Screen on the PHQ-9 Modified:

Total scores >/=11are positive

Suicidality:

Regardless of the PHQ-9 Modified total score, endorsement of serious suicidal ideation OR past suicide attempt (question 12 and 13 on the screen) should be considered a positive screen.

Interpreting the Screening Results

- Patients that score positively on the questionnaire should be evaluated by their primary care provider (PCP) to determine if the
 depression symptoms they endorsed on the screen are significant, causing impairment and/or warrant a referral to a mental health
 specialist or follow-up treatment by the PCP.
- It is recommended that the PCP inquire about suicidal thoughts and previous suicide attempts with all patients that score positive, regardless of how they answered these items on the PHQ-9 Modified.
- For patients who score negative on the PHQ-9 Modified, it is recommended that the PCP briefly review the symptoms marked as "more than half days" and "nearly every day" with the patient.
- The questionnaire indicates only the likelihood that a youth is at risk for depression or suicide; its results are not a diagnosis or a substitute for a clinical evaluation.

Depression Severity

- The overall score on the PHQ-9 Modified provides information about the severity of depression, from minimal depression to severe depression.
- The interview with the patient should focus on their answers to the screen and the specific symptoms with which they are having difficulties.
- Additional questions on the PHQ-9 Modified also explore persistent depressive disorder, impairment of depressive symptoms, recent suicide ideation and previous suicide attempts.
- Interpretation of Total Score

Total Score	Depression Severity		
1-4	Minimal depression		
5-9	Mild depression		
10-14	Moderate depression		
15-19	Moderately severe depression		
20-27	Severe depression		